THE AVIAN RESEARCH TASK FORCE

As noted above, KENETECH Windpower has been actively involved in the research of avian behavior around its Windplants since the mid-1980s. In many cases, the company has had tremendous success in reducing avian mortalities. Bird deaths, resulting from electrocution, have been significantly reduced within KENETECH's wind resource areas by retrofitting overhead electrical systems, utilizing underground wiring when feasible, and employing bird protection on all new installations.

The company has had limited success, however, determining why some birds (including certain types of raptors) are able to avoid colliding with wind turbines while other birds cannot.

In 1991, in the wake of the two-year study commissioned by the California Energy Commission, KENETECH sought the assistance of Dr. Tom J. Cade. At that time, Dr. Cade was Director of the World Center for the Birds of Prey and founding Chairman and Director of the Peregrine Fund, an organization committed to the study and preservation of raptors.

Accompanied by his colleague, Dr. William Burnham, President of the Peregrine Fund, Dr. Cade visited KENETECH Windpower's Windplants in the Altamont Pass and Solano County, California. They reviewed the research that was funded by KENETECH Windpower and the California Energy Commission's report. The initial observations of Drs. Cade and Burnham, later affirmed by the Avian Research Task Force, were as follows:

- Neither the KENETECH-commissioned research nor the study funded by the California Energy Commission was based on a sufficient number of observations to satisfy a scientific standard of verifiability. Windpower's blade painting research, for example, was not based on a statistically valid sample size to draw any conclusions regarding its viability as an effective mitigation device.
- To correct the methodological deficiency of previous studies, researchers must either increase the sample size or extend the time frame of the studies. (The reason that the number of observations in previous studies was so small is because relatively few natural encounters take place.) To address this problem, the time frame of the study could be shortened by artificially increasing the number of bird/turbine encounters through the use of trained bird populations.
- The number of recorded mortalities does not indicate *prima facie* that any of the affected populations are in danger in terms of population stability. To validate this assertion, a study should be undertaken to make a population estimate of Golden Eagles in the Altamont Pass and to calculate what proportion of Eagles observed in the Pass are migrants and which are residents.

Dr. Cade further suggested that individuals who have extensive research experience with avian physiological response to stimuli might be persuaded to design a study using trained birds. Based on his observations and recommendations, KENETECH asked Dr. Cade to assemble a task force of specialists to develop a comprehensive avian research program.

In August of 1992, the Avian Research Task Force was established to develop and implement a research program to devise effective long-term mitigation measures for KENETECH Windpower's Altamont Pass Windplant. The members of the Task force are:

- Dr. Tom Cade, Task Force Chairman and Professor *Emeritus* of Ornithology, Cornell University, Ithaca, New York. While working at the World Center for Birds of Prey and the Peregrine Fund, he pioneered adaptive breeding and reintroduction of the peregrine falcon in the wild, thus saving the species from extinction.
- Dr. Mark Fuller, National Biological Survey, U.S. Department of Interior and Director of the Raptor Research and Technical Assistance Center, Boise State University, Boise, Idaho. Dr. Fuller was formerly the chief of the Population Dynamics Group-Migratory Bird Research Branch at the U.S. Fish and Wildlife Service's Patuxent Wildlife Research Center in Patuxent, Maryland. Dr. Fuller was involved in the early use of radio telemetry in the study of avian flight and migration.
- Dr. Melvin Kreithen, Associate Professor, Department of Biological Sciences, University of Pittsburgh, Pittsburgh, Pennsylvania. Dr. Kreithen is a graduate of Cornell University and is the foremost authority on the sensory physiology of pigeons involving the modalities of sight, sound and smell. Dr. Kreithen developed of the 3-D Tracking System being employed in the field research of the Task Force.
- Dr. Vance Tucker, Professor of Zoology, Duke University, Durham, North Carolina. A graduate of UCLA, Dr. Tucker is one of the world's foremost authorities on the gliding flight of birds, particularly vultures and eagles.
- Dr. Charles Walcott, Executive Director, Laboratory of Ornithology, Cornell University, Ithaca,
 New York. Dr. Walcott, a graduate of Harvard University, is an authority on the navigation of homing pigeons.

Two research facilities were established as focal points for the Avian Research program.

MIDWAY RESEARCH CENTER

The Midway Research Center is located, on-site, at the KENETECH Windpower Windplant in the Altamont Pass. KENETECH Windpower personnel are assisting with the Avian Task Force's research on a full-time basis: Dr. David Schmitt, the resident scientist responsible for managing and conducting the Altamont-based research projects; Mr. Ron Barsic, a pigeon handler and research technician; and Ms. Denise Weingart, a wildlife response coordinator and research technician. Dr. Kreithen is the research supervisor at this location.

BOISE RESEARCH CENTER

This facility is being developed and operated under a cooperative agreement between KENETECH and Boise State University (BSU) in Boise, Idaho. At this location, laboratory-based research is underway to better understand the sensory discrimination skills of raptors. The study team is utilizing the combined facilities of BSU, the Boise Zoo, and the Raptor Research and Technical Assistance Center. The Boise staff consists of Dr. Hugh McIsaac, the resident scientist; Mr. George Carpenter, a biological technician; and Ms. Stephanie Gossett, who provides administrative support. Drs. Cade and Fuller are the research supervisors at this site.

RESEARCH STRATEGY

The first question that the research must address is: Why do some birds collide with the turbines while others do not?

- Do birds collide with turbines because of some physiological problem or is it related to behavior?
- Can birds see the turbines and, if so, how do they perceive them?
- Do the rotating blades of a turbine appear to a bird as a disk or can a bird distinguish the rotation of individual blades?
- Are the birds that collide with turbines engaged in an activity or behavior that focuses their attention away from the turbines i.e., in pursuit of prey or, perhaps, "riding" thermal wind currents and drifting backwards toward the turbines?
- Do the birds see the turbines but fail to perceive them as a threat?
- Are collisions a result of a combination of physiological and behavioral factors?
- Does the incidence of collisions vary by avian species?
- Are physiological and behavioral factors influenced by varying conditions such as the differences in lighting during the day?
- Is weather a factor in the collisions?
- Are wind speed and direction a significant factor in the collisions?
- Do certain turbine locations or alignments present more of a risk to birds than others?
- Is the size of a wind turbine blade and the rate of its rotation a relevant factor?

In order to answer these questions relating to avian sensory discrimination skills, the Task Force devised a research program to determine:

- 1. What birds can/cannot see and hear?
- 2. When, during the course of its flight, does a bird begin to take the mechanical action necessary to avoid a wind turbine?
- 3. Which colors and/or sounds do birds tend to avoid?

Through its research, the Task Force also hopes to gain a better understanding of how birds use the Altamont Pass wind energy production area (e.g., hunting, nesting, migration, etc.) in order to develop and implement effective long-term mitigation measures based on the natural behavioral traits of birds.

In order to determine what effect, if any, varying conditions have on a bird's ability to utilize its sensory perception skills, it is necessary to test and observe birds flying around the turbines at different times of the day and year, under a variety of weather conditions. Similarly, it is necessary to determine if these changes in conditions adversely affect avian activities around wind turbines.

The research of the Task Force also addresses an earlier problem — the inability to conduct tests and observations that produce statistically valid results. As previously noted, this lack of data has been a serious deficiency in every study done on this subject to date.

The Task Force research also focuses directly on the significance of avian mortalities in terms of broad population dynamics. The federal legal standard of protection for the Golden Eagle and many other species is "no losses". There is, however, a general consensus among experts familiar with this issue that this standard is not attainable. If this assertion is correct, the critical question is: What level of loss

is acceptable for each species?

In order to understand the significance of the number of mortalities reported to date, it is necessary to understand the population dynamics of the species in question. For example, what is the significance of the loss of 78 Golden Eagles over a two-year period (if it actually were to occur)?

Finally, the Task Force's research strategy also addresses the need to validate the long-term effectiveness of any mitigation action and/or device it proposes to KENETECH Windpower.

RESEARCH PROGRAM COMPONENTS

1. CONTROLLED FLIGHT EXPERIMENTS WITH PIGEONS AND MEASURED OBSERVATIONS OF WILD BIRD FLIGHT ACTIVITY

The opportunity to observe wild bird activity around wind turbines is limited by the number of encounters that naturally occur. In order to accumulate enough data to make valid observations, the Task Force elected to augment the observations of wild bird behavior in the Altamont Pass with flights of captive birds.

The objective of this research is to determine how various factors, such as a wind turbine or a string of turbines, affect the maneuverability and collision avoidance capabilities of powered or flapping flight (as well as other activities) of birds.

To carry out this research, two homing pigeon lofts have been established: one loft is located within the Altamont Pass Windplant; the other is located on the periphery of the Pass. Because the homing pigeons will return to their home loft, the utilization of these birds enables the researchers to make several hundred observations of flight behavior a week. In such flights, the homing pigeons quickly establish the direction they must take to return to their loft, making it easy to plan a release that requires the pigeons to maneuver around at least one set of turbines (usually at the release site) while enroute.

In these field tests, individual pigeons are released at various locations under a variety of conditions. Their behavior is observed and recorded visually and by the 3-D mechanical tracking device designed by Dr. Kreithen. Factors and conditions which are being tested include wind direction and velocity, lighting and range visibility, terrain, turbine type and array, and the operational status of the turbines during the test period.

In addition, field researchers will take every opportunity to observe and record the kiting (riding the wind currents) and diving flight activities of the raptors. Data from flights made under similar conditions and terrain, in areas that do not contain wind turbines, will be compiled for comparison purposes.

The findings of this research, coupled with the information obtained in Boise regarding raptor visual and auditory stimuli, will result in a series of controlled raptor flights to test the effectiveness, in terms of altered flight behavior, that occurs when the visual and/or auditory profile of a wind turbine is altered.

A critical element of flight observations is the development and utilization of the 3-D tracking system. The Task Force will be flying birds and observing natural behavior under a variety of conditions to determine what effect, if any, these factors have on the flight paths of birds maneuvering around wind turbines. The tracking system integrates several pieces of information simultaneously, enabling researchers to measure any variations in avian flight behavior.

In controlled flight situations, birds are released at scheduled intervals. The birds are then tracked by a researcher, using a siting device (similar to a telescopic sight on a rifle) which is mounted on a laser-based range finder. This tracking device is coaxially mounted on a bracket with a video camera. The bracket rests on a tripod to facilitate operation.

The system simultaneously calculates height, azimuth (direction) and distance, thus establishing a bird's location in space in relation to the turbines and the ground along the course of its flight path. With the tracking device, the Task Force can accurately trace the flight of homing pigeons around turbines and document the point at which a bird visibly reacts to the structure. During the course of experiments with controlled birds, any activity of wild birds will also be observed and recorded by the 3-D tracking system.

The tracking system is perhaps the most important means the Task Force will employ to demonstrate the effectiveness of a specific form of treatment as a mitigation measure. After compiling a database of behavior under various conditions, the Task Force will be able to quantify the differences in flight patterns, if any, that are produced by specific mitigation measures employed on or around the wind turbines.

2. SENSORY PHYSIOLOGY OF RAPTORS

In order to test the validity of employing color, patterns, shapes and/or sound as a means of alerting birds to the presence of turbines or otherwise altering their behavior, the Task Force designed a series of non-invasive experiments with raptors to establish what these birds can see or hear.

Over the years, naturalists have observed that certain colors and patterns appear to be an effective defense in startling and deflecting potential predators of a specific species. The American Kestrel, for example, has "eye spots" and a facial pattern in the plumage on the back of its head. This feature may function to ward off potential predators, or robbers of its food, that approach the Kestrel from the rear. Many predatory birds are apparently reluctant to fly directly into the face of an enemy: the "eyes" seem to be the main signal involved in this deterrence.

Three types of raptors — American Kestrels, Red-tailed Hawks, and Golden Eagles — are being studied in a laboratory setting at the Raptor Research Center in Boise, Idaho, to determine their ability to recognize wind turbines, turbine blade rotation, and the leading edge of a turbine blade. These studies will also determine which colors, patterns, and sounds are detectable to the birds and which of these produce the strongest reactions. The protocol (test procedure) for this type of experimentation is well established, having been used extensively for research on pigeons.

Data from these experiments will be used to devise field tests which will use flight situations to determine what visual or auditory stimuli can most effectively extend the range in which a bird can recognize a wind turbine as an obstacle and take evasive action.

This research effort is integrated with the controlled flight experiments taking place in the Altamont Pass. As behavioral observations are made in the field, they will be shared with Task Force researchers working on sensory discrimination in the lab in Boise.

Early research on pigeons demonstrated their ability to distinguish rotating lines. The Task Force's recent field observations of pigeon avoidance behavior around wind turbines strongly suggests that the birds are able to distinguish the rotation of individual turbine blades. Based on these observations, lab experiments will be devised to determine whether raptors are also capable of seeing rotating turbines blades in a similar fashion.

Conversely, if the sensory physiology experiments find that a particular color or pattern or sound appears to have the potential of alerting a raptor to the presence of a turbine, thereby enhancing a bird's avoidance skills, the field tests can be devised to determine the effectiveness of employing that color pattern or sound as a mitigation device in a Windplant.

It is anticipated that this type of integrated testing will begin in the Spring of 1994.

3. MORTALITY SEARCHES

At present, KENETECH Windpower field personnel are trained to spot and report any avian injury or mortality to Ms. Weingart, the wildlife response coordinator. Incident reports and any other historical data is under review by the Task Force in an effort to glean information that might suggest the cause and circumstances surrounding a bird mortality or injury.

Mortality searches, or carcass surveys, are being conducted around the Model 33M-VS turbines and a paired, or control, group of Model 56-100 turbines in the same vicinity.

This phase of research is designed, in part, to address an assertion by a few representatives of the Fish and Wildlife Service. They believe that KENETECH Windpower's new advanced wind turbine, the Model 33M-VS, presents an increased hazard to birds flying in the Altamont Pass because of its large rotor sweep area (33 meters).

The Model 33M-VS is a utility-grade wind turbine that generates electricity at rates competitive with conventional power generating fuels. The overwhelming majority of KENETECH Windpower wind turbines operating in the Altamont Pass are an earlier generation of turbines, the Model 56-100. The rotor sweep of the Model 56-100 is 59 feet.

ADDITIONAL RESEARCH

REGIONAL EAGLE POPULATION STUDY

The Fish and Wildlife Service has two distinct and often contradictory missions. The Law Enforcement division is charged with enforcing laws that are aimed at protecting individuals within a species such as the Migratory Bird Treaty Act, the Endangered Species Act, and the Bald Eagle Protection Act. The agency's other mission is to focus on population and habitat preservation, and to develop regulations that protect the vitality of a specific population.

It is unlikely that KENETECH Windpower will be able to totally eliminate the risk of avian mortalities related to wind energy production. Therefore, it is necessary to determine the amount of losses which can be sustained without jeopardizing the ability of specific species to maintain viable populations.

On September 1 and 2 of this year, KENETECH Windpower convened a conference of raptor experts which included representatives of the Fish and Wildlife Service and the Department of Fish and Game. The purpose of the gathering was to design a study program to determine the size and reproductive rate of Golden Eagles residing in the vicinity of the Altamont Pass and Solano County, California wind resource areas.

The conference participants recommended a study in which Golden Eagles will be banded and equipped with wing (patagial) tags and/or radio transmitters. A telemetry system was recommended to facilitate the ground tracking of the birds which will, in turn, enable researchers to determine the range of the age classes of resident eagles, the time eagles spend in various locales, and the type of activity occurring within these locations. It is hoped that funding support will be obtained in order to begin work in late 1993.

COLLISION IMPACT DETECTION SYSTEM

For well over a decade, KENETECH Windpower has been operating wind turbines in the Altamont Pass. Construction and maintenance personnel are on-site on a daily basis. In addition, since 1988, wildlife studies have been conducted in KENETECH's Windplants. However, it was not until August of 1993 that a bird was observed being accidently struck by a wind turbine during a pigeon flight. In an effort to capture more of these events and particularly those involving wild birds, an impact detection system is being designed for testing this Fall.

A four-camera integrated video monitoring system will be installed at a location where multiple bird-turbine collisions have occurred. These cameras will operate from dawn to dusk. The video tapes will be randomly sampled at regular intervals. If, however, a carcass is found in an area around one of the wind turbines monitored by the video camera, those tapes will be fully reviewed to see if the event was captured.

A key part of this study will involve positioning the four cameras to determine how best to capture the critical field of view. Personal observations will also be conducted as part of this study and used to compare the effectiveness of remote tracking to that of personal observations.

Given the infrequency of bird-turbine collisions and the thousands of wind turbines in the Altamont Pass, the likelihood of recording an impact is very low. Capturing a collision on video tape could enable the Task Force to fill critical gaps in its information base and possibly accelerate the timetable for making effective mitigation recommendations.

This project is an initiative of the Electric Power Research Institute (EPRI) and will be funded by EPRI and the U.S. Department of Energy (DOE) with KENETECH Windpower providing coordinating support.

SUMMARY

The research effort mounted by the Avian Research Task Force and funded by KENETECH Windpower, combined with the level of talent being marshaled to conduct the program is extraordinary. The Task Force is at a stage of testing research designs and developing equipment and field-testing procedures. Data generation is ongoing with the pace of the research expected to quicken exponentially.

A considerable effort is being made to share each step of the avian research process with interested and active parties such as environmental organizations and government agencies. This cooperative effort will ensure that the "right questions" are being asked to guarantee the objectivity and integrity of the process, of the data gathered by the Task Force, and of any mitigation measures it may recommend.

Prepared October 1993. For further information please contact Richard C. Curry, Coordinator, Avian Research Task Force KENETECH Windpower 1730 M Street, NW, Washington, DC 20036 202/833.8954 = 202/833.8960 (facsimile) = 202/434.4569 (voice mail)

^o KENETECH Windpower, October 1993.

REVIEW OF AVIAN MORTALITY DUE TO COLLISIONS WITH MANMADE STRUCTURES

Michael L. Avery
U.S. Fish and Wildlife Service
Ann Arbor, Michigan

INTRODUCTION

Birds have undoubtedly been colliding with manmade structures ever since humans started building things skyward. The earliest documented instances of collision mortality in this country are from the late 1800's (Coues 1876, Merriam 1885) and the problems continue unabated today. The total axian mortality due to collisions with manmade obstacles is probably greater every year as buildings, towers, chimneys, overhead power lines, and other structures are erected in ever-increasing numbers. Weir (1976) provides an excellent review of bird migration, weather, and collision mortality associated with various types of structures.

The biological significance of collision mortality to any species is unknown. There are too many other factors, such as total population size, natural mortality levels, and other human related influences, for which insufficient data exist to be able to put the collision mortality factor in proper perspective. Recognizing these limitations, Banks (1979) recently analyzed axian mortality, including collisions, related to human activities and estimated that about 4.75 million (0.05%) of the 10 billion annual bird deaths result from collisions with manmade structures.

In this paper, I will discuss each of the major sources of avian collision mortality and briefly review any pertinent investigations recently completed, currently in progress, or planned for the near future. Where relevant, comparisons with mortality estimates of Banks (1979) and others will be made. In addition, measures that have been developed to reduce the amount of various types of collision mortality will be mentioned.

Basically, this is a review, and the information presented herein was obtained from the published literature and through personal communications with investigators in this field. Most of the available reports and articles were previously compiled in an annotated bibliography (Avery et al. 1978a). For some of the mortality categories, certain analyses were made to obtain an indication of the magnitude of the annual losses attributable to these sources. These analyses are described in the appropriate sections of the discussion.

DISCUSSION OF MORTALITY SOURCES

TV and Radio Towers

Since the mid 1950's, bird losses at TV and radio towers have attracted much attention, especially throughout the Eastern United States. These structures and their supporting guy wires comprise the largest single category among the available reports and articles dealing with collision mortality (Avery et al. 1978a). The annual losses due to collisions with towers have been estimated at 1.0 to 1.25 million birds (Aldrich et al. 1966, Banks. 1979), which represents approximately 0.013 to 0.016 percent of the total estimated annual avain mortality (Mayfield 1967, Banks. 1979).

Because mortality can vary considerably from year to year in deriving my own estimate of the yearly number of tower-killed birds, I considered only studies of at least 10 years' duration. Just four studies meet this criterion, three of these report fail mortality only, so I restricted the analysis to fall losses. Table 1 summarizes the fall mortality data at the four locations, each tower has been under investigation for at least 11 years. The wide range in fall mortality figures underscores the necessity for long-term investigations. On a per tower basis, WCTV at Tallahassee has recorded the highest seasonal mortality (1305), followed closely by WSGN at Nashville (1155). The lowest seasonal mean is from WIVB in Buffalo (61).

ort bird losses due to striking system of telegraph wires as a along one 3 mile (5 km) stretch, ines criss-crossing the country miles (853,300 km) of electric ic Institute 1978), over 333,000 ow 22 kV (U.S. Department of m) of aerial telephone cable and imission line circuit may consist one or two ground wires. Thus, 1 overhead wires. A structure km are strung. A structure km may uctor or wire-km.)

ne unit "wire-km" is used in the nventional "circuit km" because viguestion than is the number of ter of circuit km by four gives an invert structure km to wire-km due to striking overhead wires, lurce. Only those studies design-dimost of these were unsuitable along the number of conductors. As a result, data from only a few arized in Table 2.

ns involve power lines (above 22 biased. Nevertheless, the data 30 kV power lines examined by These two study sites were the mat the mortality rates there are

the variation in the duration of fall and assigned 0.5 yr. to any sty-June) or fall (July-December). te (Table 2) was examined for a lee 1978). 22 October 1977-31 1978). In my calculations these Table 2) Table 2 lists the actual these.

refestimate the actual mortality me periods used in the calculating only the actual numbers of rates. Loss totals corrected for pecause not all of the studies.

carricularly great in view of the indicates that an intermediate in characterizing bird losses at abitats having comparable birdires differences in habitat and mortality may exist even between projections of total mortality caution. Lee (1978) and Meyer of flight intensity to collision mortality catterier habital types it would paper to nationwide or regional

published the proceedings of a with electric power transmission describer currently underway or sees due to collisions with power intinuing to conduct what is to alty and behavior at power tines. Tusty attempt to account for the ger removal of carcasses and in

vestigator error. Also, they have acquired considerable data on bird flight reactions near power lines and have characterized collision rates in terms of the flight intensities across the lines.

Second, the U.S. Fish and Wildlife Service (FWS) is currently involved in three proiects in the Upper Midwest. At Red Wing, Minnesota, the Service, in cooperation with Northern States Power Company, is investigating the effects of a proposed 345 kV power line crossing the Mississippi River. Preconstruction studies include radar monitoring of bird movements through the area and laser beam simulation of a power line to determine the potential bird strike incidence. Postconstruction efforts will concentrate on assessments of bird mortality at the line. In North Dakota, the FWS area of fice at Bismarck initiated a post card survey and questionnaire on bird strikes at overhead wires to identify specific problem areas in the state. The survey elicited 37 responses concerning 66 losses in 1978, and the survey will continue at least through 1979 (Cernohous 1978). The Northern Prairie Wildlife Research Center (FWS) in Jamestown, North Dakota, in cooperation with several utility companies, has designed a study to determine the magnitude and significance of bird strike mortality to avian populations. This study, which has not yet commenced, will concentrate first on known trouble spots, and then expand to other areas in the state if the level of mortality in dicates a second phase is warranted. The mortality due to collisions with wires will be evaluated in the context of other, documented mortality factors such as disease, predation, and hunting

Finally, the Electric Power Research Institute, Palo Alto, California, has contracted for a multi-year study of bird flight interactions with overhead transmission lines. The initial phase of the study, expected to begin in January 1980, will be concerned with the development of techniques and procedures for data collection and analysis of bird interactions with transmission lines.

These and, hopefully, other studies will ultimately provid answers to the questions concerning the magnitude and biological importance of bird iosses at overhead lines, and will provide necessary direction for reducing the problem

CONCLUSION

From this review, it is apparent that the primary source of collision mortality among birds are not the spectacular, episodic events recorded at stuctures such as TV towers and power plant stacks, but are the small, incremental losses associated with the millions of kilometers of power and communication lines and the billions of grass win dows throughout the country. Losses from the latter two sources are difficult to document because mortality at any one site is usually so small it goes unnoticed. However windows and overhead wires together with road-related losses (Banks 1979), may account for hundreds of millions of bird deaths annually. Whether or not such mortality is of any biological significance remains to be determined, but in view of the deleter busie fects of policition, habital destruction, and other humanine aled activities, the impacts of collision mortality cannot be ignored.

ACKNOWLEDGEMENTS

I am grateful to the following persons for their comments on an earlier version of this manuscript. Kenneth D. Hoover, Jack M. Lee, Jr., James Lewis, P. Kent Schreiber, and Paul F. Springer. Arthur R. Clark kindly provided unpublished data from his tower kinstudy in Buffato. New York, At the National Power Plant Team, the efforts of Carol Duren, who ably typed several drafts of this paper, and Kathy Gilden, who researched some of the information contained herein, are greatly appreciated.

LITERATURE CITED

- Aldrich J.W. R.R. Graber, D.A. Munro, G.J. Wailabe, G.C. West, and V.H. Canalane, 1966, Report of Committee on Bird Protection, Auk 83, 465-467.
- Andersen Harild P. and D. Bloch. 1973. Birds killed by overhead wires on some locations in Denmark. Dan. Ornithol. Foren. Tidsskr. 67:15:23.
- Avery M. and T. Ciement. 1972. Bird mortality at four towers in eastern North Dakota tall 1972. Prairie Nat. 4-87-95.
- Avery M.L. P.F. Springer, and J.F. Cassel. 1975. The effects of a tail tower or nocturnal bird migration. a portable ceilometer study. Auk 93,281,291.

- Avery Mill P.F. Springer and N.S. Davey 1978a. Avian mortality at manimal positive tures, an annotated bibliography, U.S. Fish Wildl. Serv. FWS/CBS 15/83, 108 pp. (Supt. Docs. Stock No. 024-010-00472-0)
- Avery M.L. P.F. Springer, and J.F. Cassel 1978b. The composition and seasonal variation of bird losses at a tall tower in southeastern North Dakora. Am. Birds 32 1114-1121
- Baird J. 1970. Mortality of fail migrants at the Boyiston television tower in 1970. Chickadee 40 17-21
- 1971 Mortality of birds at the Boylston television tower in Sectember of 1971 Chickadee 41 20-24
- Bardwin, D.H. 1965. Enquiry into the mass mortality of nocturnal migrants in Chrano. Final report Ont Nat 3.3-11
- Banks, R.C. 1979. Human related mortality of birds in the United States, U.S. Fish Wildli Serv Spec Sc. Rep Wild: 215 16 pp
- Barrold, W. 1976, Bird kill at a TV tower, Redstart 43,109
- Bier v. M.L. 1973, 1971, fall television tower casualties in Nashville, Migrant 44,5,6
- Bradstreet MISIWI, and PISI Woodford in di Nocturna imigration, upng Point Bird Observatory Ten Year Report 60169 pp. 19/23
- Broughton, D. 1977. The bird kill problem at Ontario Hydro's thermal generating staons, a study of nocturnal migrant mortality due to limpingement at lighted stacks. Ontario Hydro CTS-07017-1 30 pp.
- Casset U.F. D.W. Kiel J.J. Knodel, and J.M. Wiehe. 1979. Relation of birds to certain botwer transmission lines in central North Dakota, Zool, Dept. N. Dak, State Environment -3:32
- Cernonous L. 1978 Untitled U.S. Fish W. dt. Serv. Area Off. Bismarck, N.D. Unbubl. res 6 pp
- Scottan WW and RR Graber 1958 Affrantion of nocturnal migrants by ignision a elevision tower. Woson Buil 70 316 380
- Court Eliters The destruction of birds by telegraph with Amilian in 194 19-
- Crawford Pig. 1914, Bird casualities at a Leon County, Florida Tvirower, Circ cur 1946. September, 1913, Tar. Timbers Res. Stn. Bust. 18, 21 pp.
 - 1978. Autumn bird basualties at a northwelt Fig. 1997, 1994-1981, 1985. A 1,5 Bu 90 035 346
- Epiton Electric institute in 978. Statistical vearbook of the Electric Lincoln housing harm . :::
- Fire LD and TW French 1971 Birds in downtown Atlanta (1971) Chole
- French T W 1973 Notes from Atlanta (1971) and 1972 Once 38 39 43
- Garler, A.F. 1962. Bird casualtes at a Nashville TV tower, Migrant 33,58,50
- Noodbabrure IKIA (1974a) Fail 1972 relevision tower basuar As in Nashville IMIçlanti
- 📖 = 19740 Fall 1973 television tower casualties in Nashville iMigrant 45 51 59 19% Nachville lelevición tower casualt esi 19%. Migrant 46,49 5%

Howard 16.2	W 1	1963	Migr	at on
	1966	~ 2.50	ver ki	j; 1
	1967	Tower	kill 1	967
	1968	Tower	K II 1	968
	1969	Migra	con c	asua
	1970	1970	lower	kal
	1971	Tower	4 (1)	197
	1972	Towe:	Kill .	197
cage	·973	ASYE seri	- TV 3	asua
	1974	Tower	asu	aitie
	-977	,:vē.	'k 1	197
Jackson ing d	w B sontra	M Te ct sen	ان عاد فستسو	anual
Jarosio# cedu	v 8 N ires F	∙arg resent	A re ed at	liew the
Johanse Onta	r К ′СН,	1975 droige	₽ø, e nera	w af
Johnson Masi	ningto	and in Stati	3 E e :We	Hu.
Johnsto Aur				
Kibbe D) = •	976 7	.ė ,9.	~ ;
Kiem D Spur	Je. J	· 479 - 75 3	સુ ૧૩/ ૫૧/	gv o
.şr.e. 1. f				
 -	1955	, ڊ د*	s on t	'sv.e
		.دِ دِ		
		3.		

1976 Nashul e le evis

1958 1959 1950 Migrant 3

33 7 8

34 65 56

1962 Migration data

📖 1963a Caslares ar 🤉 . 1963b Morrally of

** an mortal ty at man made struct

Serv. FWS:085 18 58 106 pp.

The composition and seasonal varial Hastern North Dakotal Am Birds

Boviston relevision rower in 1970

-. sign tower in September of 1971

🦥 🖟 **ct noct**urnal **m** prantiti in Chitano

the United States (1.5) First Wild

40.00

ries ni**Nash.** je Mizram 44.5.6 Riji Pruma imigranom upng Point Biro

ेट Mydrois thermal ceneraling sta अ**ाट impingeme**nt ar i ghted stacks

* 1979 Relation of birds to certain is Zool Dept. N. Dak State Univ.

Area Off Bismarck N.D. unpub

ti nocturna imigranto più istito no a

17 AM AT NEW 17 134 735

ty Florida Tyroxie - Childen 1965 - 27 bb

west Florida TV tower 1973 1975

thine Electrician is industry, 1977

grider Atama ira 157 Cice

""+"& Orde 38 344

g tower Migram 39.55 ().

· lessane in the color this con-

in Nashiyi el Mişları 46 (1954) 1974 - Mistarti ek 2006

1975. Nashville felevision fower casualties, 1975. Migrant 47-8-10. Howard, Willinger Migratinh Lersus progress, But Chemung Valley Audubon Soc 1966 TV tower kill 1966 Bull Chemung Valley Audubon Soc 191 1967 Tower kill 1967. Bull. Chemping Valley Audubon Soc. 20.2 _ 1958 Tower kill 1966 Bull Chemung Valley Audubon Soc. 21.2 1969 Migration casualities 1969 Bull Chemung Valley Audubon Soc 22.3 ____ 1970 1970 towerk Bur Chemong Vall, Audubor Soc 23.1 ____ 1971 Tower kii: 1971 Bull Chemung Valley Audubon Soc 24.1 1972 Tower kill 1972 But Chemung Valley Audubon Soc 25 1 2 _ 1973_WSYETV casuartes 1973_Bull Chemung Valley Audubon Soci 26 (no ji başe numberi 1974 Tower casuarties 1974 But Chemung Valley Audubon Soc. 27.4 _ 1977 Tower kill 1977 But Chemung Valley Audubon Soc 30 2/3 Jackson W.B. M. Temme, and W.A. Peterman, 1977. Davis Besse pird hazard monitor. ing contract, sem annual report, January 1977, Toledo Edison Co. 21 pp. Jaroslow, Bit., 1979. A review of factors involved in bird tower kills, and mitigative procedures. Presented at the Mitigation Symposium, 16.20 July. Ft. Collins. Colorado. Johansen K. 1975. Review and analysis of bird impingement and stack illumination at Ontario Hydro generating stations, Ontario Hydro, Rep. 75073, 42 pp. Johnson RE and GE Hudson 1976 Bird mortality at a glassed-in walkway in Washington State, West, Birds 7,99 107 Jannesten C. A. and T.P. Haines. 1957. Analysis of massit indimortality in October, 1954. 43. "4 44" 458 Kibbe 1 F 1976. The tall migration Niagara Champtain region. Am. Birds 30 64-66. Kiem, D. 1919, Biology of collisions between birds and grass, Ph.D. Dissertation Scuttern tind's Univ. Carbondale Laskey: A.R. 1951. Another disaster to migrating birds at the Nashville airport. Migrant 22 57 60 _ 1956. Television towers and nocturnal bird migration. Migrant 27 66-67. _ 1957 Television tower casualties. Nashville: Migrant 28:54:56. 1980. Bid migration casualties and weather conditions autumns 1958 1959 1960 Migrant 31 61-65

1962. Migration data from television tower casualties at Nashville. Migrant

1963b. Morrally of hight migrants at Nashville TV towers, 1963. Migrant

____1 19-3a. Casualtes at WS/x TV tower in autumn 1960. Migrant 34-15.

37 76

34 6: ...

	1964	Data	!rom	tre	Nashville	ΤV	owe.	casualties	au'umr	 M grant	
35 95	96										

- _____ 1965. Autumn 1965 TV tower casualties at Nashville. Migrant 36 80 81
- _ 1966 TV tower casualties at Nashville spring and fall 1966 Migrant 37 61-62
- ___ 1968. Television tower casualties at Nashville, autumn 1967. Migrant 39,25-26.
- 1969a TV tower casualties at Nashville in autumn 1968. Migrant 40 25-27
- _ 1969b. Autumn 1969 TV tower casualties at Nashville. Migrant 40.79/80.
- 1971. TV tower casualties at Nashville, spring and autumn, 1970. Migrant 42 15 16
- Lee J.M. Jr. 1978. Effects of transmission lines on bird flights, studies of Bonneville Power Administration lines, Pages 53 68 in Mig. Avery, ed. Impacts of transmission thes on birds in flight U.S. Fish Ailal Serv FAGOBS:78:48
- LGL Limited (1972) An examination of the bird impact problem at the Nant coke plant of the Ontario Hydrorelectric system, phase it autumn, 1972. Unpublined, 32 pp.
- Mai/ erd H 1967 Shed few tears Audubon Mag 69.61.65
- McGrauphin, D.C. 1977. Description and chronology of events on Shake Creek embank ment, U.S. Fish Wild: Serv. Audubon Wildlife Refuge. Colenarbor, N.D. Unpubl. rep.
- McKenna, M.G., and G.E. Aliard, 1976. Avian morrality from wire collisions, N. Dak, Out. doors 39(5) 16:18
- Merciam, C.H. 1985. Presiminary report of the committee on bird migration, Aux 2,53,57
- Meyer, u.R. 1978. Effects of transmission lines on tind from behavior and collision more tality Bonney le Power Admini Portland Ore 200 pc.
- Stout 100 and G.W. Cornwell 1976. Nonrunting mortality of fleaged No. to American Agrerton a N d. Mamh 40 681 693
- Tenner U.T. 1964, Bird mortains during night migrardh. Cotober 1964, Migrant
- THIBEDTONY Publishing Corporation (1918) Telephony (Chicago)
- Tummye Miliw Bi Jackson, and wilk Pererman (1914) Daylik Benselo, dinase dimontro. of Dustriator annual report variable (91%) folles emplo Stocklos
- 3.6. Debaitment of Energy (1914) Statistic lights with your equiversity of the control States (1911). Energy, information Agric 1943, 402.
- USI Fluttland Wild ite Service (1978) impacts of transmission lines on birds in highriphs the sings of la Workshop (FMS CBS 1848) 181 bb (Subi) Cots (Brock No. 144) 110 (148) 8
- ${\it Mex} (B,C)$ 1913a. Bird wills at the Lennox plant of the Ontario hydroelectric system, spr nt 1913 Blue B 2013 24
 - 1970b. Bird wis at the Lernox generating blant autumn 1913. Blue Bir
 - THIS Armitiated bib objects of bids, is at manimate springles, a review of the varient and solutions. Can take 3 Servicion Regions was 85 pc.
 - 1971. Blook it styffe cent ik generating waron, spring and autumn 1971. Blue 8 14:542

	Falletin see	Nash	Nashville'		Bullalo ³	
	<u> </u>	WS.W	WSIX	WCH	WKBW	WIVB
(m) bounded about more	****	415	285	591	326	321
Control of the second s		2	2	Ξ	22	15
Transfer of Buck collected	- -	H. 18.	6224	5389	4117	977
Marine Or Dings Constitution		1.61	311	4,30	360	1.9
Magazia	1/62 01	56 3683	56 1553	24.1824	2151 91	1 190

255 WSYE

Elmira.

1778 36/4

> 1965 Gamer 1971 1969 1969 1:368 1965 ÷ ± -14, 11 1960 185hry 1956 1957

1978

1 13W Ord 1974

Table 1. Fall lower kill data from local location, in the earliern United States

1

1 67

"

	Tallahas are o − c WCTV	Na	Stiville?		Bullalo*			
		WSM	WSIX	WGR	WKBW	WIVE	WSYE	
Towns height above ground (m)	ROF	415	285	291	326	321	255	
Fall stations monitored	,11	16	20	11	12	12	14	
Namber of birds collected	27.1(n)	18568	6224	5389	4317	726	8771	
Mean fall fell	1.05	1161	311	490	360	61	627	
Harage	2.16 397.1	56 3683	56 1553	24 1824	16 1312	1 190	45 3874	

G JC 84

ci

3

0

23

Ş

3": Denavici and

.0.5

3

protrematine Name

.ec 32 pp.

To fights, studies of Bonney Re

A-11-1-SEA

Migran:

TG and autumn

1967 Migrant 39.25

and tall 1966. Migrant

37

r)

Migrant 36 80

<u>ج</u> بن

*Crawford 1974 1978

œ

Zkarkey 1956, 1957, 1960, 1962, 1963a, 1963b, 1964, 1965, 1966, 1968, 1969a, 1969b, 1971, Ganier 1962, Bierly 1973, Goodpasture 1974a, 1974b, 1975, 1976

A R. Clark. Buffalo Museum of Science. Buffalo. N.Y. Personal communication

Howard 1963, 1966, 1967, 1968, 1969, 1970, 1971, 1972, 1973, 1974, 1977, Kibbe 1976

Table 2 Rates of bird loss due to collisions with power lines

	Number of	,	_	
	dead birds		Study	Losses/wire
Location	collected	Wire kin	period	km/yr
Denmark!	105	432.3	19 da (0.5 yr.)	0 49
Oregon/Washington ^a			• • • •	
Lower Crab Creek	12	112	6 mo (1 yr)	0.91
Frenchman Hills	?	8.8	3 mo (0.5 yr.)	0.45
Rocky Ford Creek	U	196	3 mo (0 5 yr)	0
Bybee Lake	76	14.0	9 mo (1.5 yr.)	3 62
Riekkola	1	24	3 mo (0.5 yr.)	0.83
Rear River	0	2.4	3 mo (0.5 yr.)	0
North Dakota			, ,	•
Snake Creek!	632	50.4	24 mo (5 yr.)	2 5 1
Underwood At	49	25.8	6 mo (1 5 yr)	1 27
Underwood 81.1	4, 1	435	4 mo (1 yr)	1 17

'Andersen Harild and Bloch (1973)

*Lee (1978) Meyer (1978)

McKenna and Allard (1976). Michauchlin (1977). Cernohous (1978).

*Carsel et al. (1979).

Collision Course:

The Hazards of Lighted Structures and Windows to Migrating Birds

A Special Report by Lesley J. Evans Ogden for World Wildlife Fund Canada
and the Fatal Light Awareness Program, September 1996

APPENDIX 1, Bird Collision Literature Summary Table (TV Tower Collisions)

Locations	Years	No. Killed		Predominant Species/Groups	Reference *
WJBF-TV, Aiken, SC, USA	1962	400		Red eyed Vireo	766
Alleman, Iowa, USA	1972	726		406 (40%) Warbler	420
Baltimore, MD, USA	1964	1032	37	300 (29%) Ovenbird	669
Barrie, ON, CAN	1974	4900		1000 (20%) Bay-breasted Warbler, 900 (18%) Ovenbird	337
Boston, MA, USA	1958	300		Warbler, Vireo	63
Boyleston, MA, USA	1971	158	29	134 (85%) Warbler, 95 (60%) Blackpoll Warbler	62
Boylston, MA, USA	1970	350	29	266 (76%) Warbler	61
Buffalo, NY, USA	1978	359	51	44 (15%) Blackpoll Warbler, 36 (10%) Ovenbird, 35 (10%) Swainson's Thrush, 25 (7%) Red-eyed Vireo	892
Buffalo, NY, USA	1974	651		Warbler	169
Buffalo, NY, USA	1970	534	46	105 (20%) Yellow-rumped Warbler, 63 (12%) Black-throated Blue Warbler	775
Carolinas, USA	1962	4189	61	American Redstart, Ovenbird, Vireo	5
Cedar Rapids, IA, USA	1963			Thrush, Warbler	585
Chapel Hill, NC, USA	1956	2500	40	Warbler, Thrush	Chat (1957) Mar
Chapel Hill, NC, USA	1956	2500		Warbler	159
Charleston, NC, USA	1954	1000- E18	24	Warbler, Common Yellowthroat	Chat (1954) 18(4)
Charleston, SC, USA	1962		Į.	Red-eyed Vireo, Ovenbird, American Redstart	766
CHRE-TV, Regina, SK, CAN	1965	172		Warbler	90
CKCK-TV, Regina, SK, CAN	1965	227		Warbler	90
CKVR-TV, Barrie, ON, CAN	1975	175		Bay-breasted Warbler, Ovenbird, 414 (10%) Red-eyed Vireo, 313 (8%) Chestnut-sided Warbler	840
Columbia, MN, USA	1963	941		Red-eyed Vireo, Ovenbird	585
Dallas TX, USA	1960	11	1	Yellow Rail	85
Davenport, IA, USA	1960	281	25	Thrush, Warbler	506
Dayton, OH, USA	1966	305	49	Red-eyed Vireo, Golden-crowned Kinglet, Ovenbird	590
Des Moines, IA, USA	1974	1500		750 (50%) Red-eyed Vireo	415
WEAU-TV, Eau Clair, WI, USA	1957	1525	40	Warbler	404
WEAU-TV, Eau Clair, WI, USA	1957	2972	42	Warbler	116
Elmira, NY, USA	1966	270		Ovenbird	644
Elmira, NY, USA	1969	300		Bay-breasted Warbler	647
Elmira, NY, USA	1972	540	55	Warbler	649
Elmira, NY, USA	1975	800	40	198 (25%) Bay-breasted Warbler, 78 (9.8%) Ovenbird, 110 (14%) Magnolia Warbler	411

Erie County, NY, USA	1977	1397	50	168 (12%) Bay-breasted Warbler, 154 (11%) Ovenbird, 112 (8%) Magnolia Warbler	172
FL, USA	1964	4707	37	4646 (99%) Warbler	154
FL, USA	1971	2500	42	Warbler	394
FL, USA	1972	1347	49	1199 (89%) Warbler	744
Grand Bahama Is, USA	1966	136	22	Gray-cheeked Thrush, Blackpoll Warbler	401
Jacksonville, FL, USA	1964	2000		1900 (95%) Warbler, 273 (14%) Blackpoll Warbler	193
Jacksonville, USA	1970	146		Warbler	633
Jacksonville, USA	1967	174		Ovenbird	635
KCMO-TV, KS, MO, USA	1975	67		23 (34%) Mourning Dove	289
KOMU-TV, Columbia, MO, USA	1954	1887	63	354 (19%) Common Yellowthroats, 313 (17%) Gray Catbird	290
KROC-TV, Ostrander, MN, USA	1961- 62, 1972- 74	3507	84	619 (18%) Northern Waterthrush, 516(15%) Red-eyed Vireo	729
KTOL-TV, Coweta, OK, USA	1974	117	28	64 (55%) Nashville Warbler	554
Lawrence, KS, USA	1969	19		Thrush and Sparrow	545
Lennox Power Plant, Barrie, London TV ON, CAN	1974	7550		1359 (18%) Bay-breasted Warbler, 1129 (15%) Red-eyed Vireo, 1038 (14%) Ovenbird, 920 (12%) Magnolia Warbler	283
London, ON, CAN	1970			Ovenbird, Warbler	279
Madison, WI, USA	1968	493	33	Thrush, Warbler, Warbler	677
Maryville, MO, USA	1972	71	33	Sparrow	69
MI, USA	1954	230		Blackpoll Warbler	224
Moosejaw, SK, CAN	1959	33	13	Yellow-rumped Warbler, Orange-crowned Warbler	426
NY, USA	1959	110		Warbler, Vireo	663
Olney, IL, USA	1978	622	36	498 (80%) Warbler	351
Omega Tower, LaMoure, ND, USA	1973	1417	51	Finch	38
Omega Tower, LaMoure, ND, USA	1972	255	58	Finch	39
Omega Tower, LaMoure, ND, USA	1972	226	66	Warbler	39
Omega Tower, LaMoure, ND, USA	1971- 73	937	102	Warbler, Vireo	42
Omega Tower, LaMoure, ND, USA	1971	152	41	Warbler, Vireo	25
ON, CAN	1961	1115	57	156 (14%) Ovenbird, 99 (8.9%) Chestnut- sided Warbler, 91(8.2%) Bay-breasted Warbler, 91 (8.2%) Red-eyed Vireo	64
ON, CAN	1962	3446	66	Thrush, Warbler, Vireo	65
ON, CAN	1963	1190	71	Thrush, Warbler, Vireo	66
Orion, IL, USA	1959	88		Swainson's Thrush, Warbler	505
Philadelphia, USA	1948			Warbler	603
S. Atlantic coast, USA	1954			Ovenbird, Red-eyed Vireo	156
S. Erie County, NY, USA	1967- 71	4094	82	450 (11%) Ovenbird, 409 (10%) Golden Crowned Kinglet, 287 (7%) Blackpoll Warbler, 287 (7%) Gray-cheeked Thrush, 246 (6%) Vireo	167
South Bend IN, USA	1959	49		Swainson's Thrush, Warbler	505
Topeka Tower, KS, USA	1967	800	43	240 (30%) Nashville Warbler	152
Topeka, KS, USA	1955	16	2	15 (94%) Mourning Warbler, 1 (6%) Connecticut Warbler	83

Various	1	16118		2498 (15.5%) Ovenbird, 1950 (12.1%)	259
				Tennessee Warbler, 1418 (8.8%) Red-eyed	
		l 		Vireo, 1418 (8.8%) Magnolia Warbler	
Vero Beach, USA	1970	31		Warbler	633
WBAL-TV, Baltimore, MD, USA	1970	1965	43	489 (25%) Ovenbird, 410 (21%) Red-eyed Vireo	671
WBAL-TV, Baltimore, MD, USA	1973	180		Warbler	673
WBAL-TV, Baltimore, MD, USA	1970	1800	41	435 (24%) Ovenbird, 391 (22%) Red-eyed Vireo, 148 (8%) Black and White Warbler, 115 (6%) Common Yellowthroat, 81 (5%) Magnolia Warbler	
WBAL-TV, Baltimore, MD, USA	1964- 66	3595	74	899 (25%) Ovenbird, 468 (13%) Black-and- white Warbler, 395 (11%) Magnolia Warbler	136
WBDO-TV, Orlando, FL, USA	1970	2790	51	Warbler	633
WCIX-TV, Homestead, USA	1970	300		Warbler	633
WCSH-TV, Sebago, USA	1973	300		Warbler, Thrush	292
WCTU-TV,Tallahassee, USA	1962	249		Red-eyed Vireo	578
WCTV-TV, Leon County, FL, USA	1963	735		81 (11%) Bobolink	191
WCTV-TV, Leon County, FL, USA	1964	709		335 (47%) Yellow-rumped Warbler	713
WCTV-TV, Leon County, FL, USA	1973- 75	3864	109	896 (23%) Red-eyed Vireo, 219 (6%) Ovenbird, 159 (4%) Common Yellowthroat, 140 (4%) Magnolia Warbler	899
WCTV-TV, Tallahassee, USA	1960	237	53	Warbler	633
WCTV-TV, Tallahassee, USA	1960	384		230 (60%) Sparrow	637
WEAU-TV, Eau Clair, WI, USA	1968	145		Kinglet, Warbler	629
WECT & WWAY-TV, SE NC, USA	1971- 77	7270		1023 (14%) Common Yellowthroat, 925 (13%) American Redstart, 865 (12%) Ovenbird, 701 (10%) Red-eyed Vireo, 549 (8%) Black-and-white Warbler	888
WECT-TV, NC, USA	1971- 72	3070	84	Warbler, Sparrow, Thrush, Vireo, 583 (19%) Common Yellowthroat, 288 (9.4%) Black- throated Blue Warbler, 267 (8.7%) Ovenbird, 218 (7.1%) Yellow-rumped Warbler, 163 (5.3%) Gray Catbird	Chat (1976) 140(1)
WEHN-TV, Deerfield, NH, USA	1959	130		74 (57%) Ruby-crowned Kinglet	661
West Brands, IA, USA	1970	58	16	Kinglet, 14 (24%) Nashville Warbler, 9 (16%) Ruby-crowned Kinglet, 8 (14%) Yellow-rumped Warbler, 7 (12%) Golden-crowned Kinglet	1022
WFMJ-TV, Youngstown, OH, USA	1975	1057	39	Warbler, 317 (30%) Ovenbird	78
WFMS-TV, Youngstown, OH, USA	1977	315		Bay-breasted Warbler, Blackpoll Warbler	873
WHEN-TV, Syracuse, NY, USA	1959	45		Thrush, Vireo, Warbler	662
WHIO-TV, Dayton, OH, USA	1967	348	45	Red-eyed Vireo, Warbler	591
WHNT-TV, Huntsville, USA	1976	42	18	27 (64%) Warbler	896
WIS-TV, Columbia, SC, USA	1969	500	20	Warbler, Thrush, Vireo, Common Yellowthroat, Magnolia Warbler	165
WJBF-TV, Aiken, SC, USA	1962	200	32	48 (24%) Swainson's Thrush	Chat (1963), Mar
WJBF-TV, Aiken, SC, USA	1962	400	32	239 (60%) Red-eyed Vireo	601
WMC-TV, Memphis, TN, USA	1961	19	11	Warbler, Vireo	176
WMC-TV, Memphis, TN, USA	1964	99	21	58 (58%) Red-eyed Vireo	176
WPSK-TV, Clearfield Co. PA, USA	1969	75		Brown Creeper, Kinglet, Warbler	1039

WSM & WNGE-TV, Nashville TN, USA	1976	406	43	63 (16%) Ovenbird, 61 (15%) Tennessee Warbler, Magnolia Warbler, Bay-breasted Warbler	920
WSM & WSIX-TV, Nashville TN, USA	1971	3560		Warbler, 845 (24%) Tennessee Warbler, 631 (18%) Ovenbird, 429 (12%) Black-and-white Warbler, 420 (12%) Magnolia Warbler	452
WSM-TV, Nashville TN, USA	1967	160	12	115 (72%) Blackpoll Warbler	448
WSM-TV, Nashville TN, USA	1968	5408		4380 (81%) Warbler	450
WSYE-TV, Elmira, NY, USA	1963	200	36	Warbler	342
WSYE-TV, Elmira, NY, USA	1968	260	30	Warbler	346
WSYE-TV, Elmira, NY, USA	1973	465	39	Warbler	351
WSYE-TV, Elmira, NY, USA	1974	844		246 (29%) Bay-breasted Warbler	352
WSYE-TV, Elmira, NY, USA	1977	3874	48	1227 (32%) Bay-breasted Warbler, Magnolia Warbler, 311 (8%)Ovenbird, 218 (6%) Swainson's Thrush	353
Youngstown, OH, USA	1975	1050		305 (29%) Ovenbird	27

Bold indicates where number given is an estimate or a minimum
*Numbered references refer to Avery, M.L., P.F. Springer, and N.S. Dailey (1980). Avian mortality at man-made structures: An annotated bibliography (revised). U.S. Fish and Wildlife Service, Biological Services Program, National Power Plant Team, FWS/OBS-80/54. 152pp.